

The Star Formation History and Stellar Assembly of High Redshift Galaxies

Casey Papovich

Spitzer Fellow, Steward Observatory, 933 North Cherry Avenue, Tucson, Arizona, 85721, USA
 email: papovich@as.arizona.edu

Abstract. I discuss current observational constraints on the star-formation and stellar-assembly histories of galaxies at high redshifts. The data on massive galaxies at $z < 1$ implies that their stellar populations formed at $z > 2$, and that their morphological configuration was in place soon thereafter. *Spitzer* Space Telescope 24 μm observations indicate that a substantial fraction of massive galaxies at $z \sim 1.5 - 3$ have high IR luminosities, suggesting they are rapidly forming stars, accreting material onto supermassive black holes, or both. I compare how observations of these IR-active phases in the histories of massive galaxies constrain current galaxy-formation models.

Keywords. galaxies: evolution, galaxies: formation, galaxies: high-redshift, infrared: galaxies

Most ($\sim 50\%$) of the stellar mass in galaxies today formed during the short time between $z \sim 3$ and 1 (e.g., Dickinson et al. 2003, Rudnick et al. 2006). Much of this stellar mass density resides in massive galaxies, which appear at epochs prior to $z \sim 1-2$ (see, e.g., McCarthy 2004, Renzini 2006). The fashionable scenario is that galaxies “downsize”, with massive galaxies forming most of their stars at early cosmological times, with less-massive galaxies continuing to form stars to the present (e.g., Juneau et al. 2005). However, it is still unclear when and where the stars in these galaxies formed. For example, it may be that stars form predominantly in low-mass galaxies at high redshifts, which then assemble over time to form large, present-day massive galaxies (e.g., Kauffmann & Charlot 1998).

At $z \lesssim 1$ massive galaxies exist on a fairly prominent red sequence (e.g., Blanton et al. 2003, Bell et al. 2004, Willmer et al. 2006), are largely devoid of star formation and evolve passively. Forming such a red sequence is a challenge for contemporary hierarchical galaxy formation models (e.g., Davé et al. 2005) without including some agent to suppress star formation (the favorable mechanism is feedback from AGN; e.g., Croton et al. 2006, Hopkins et al. 2006). In hierarchical models, the most massive galaxies continue to grow via satellite accretion. To maintain the red sequence this growth must occur without continued star-formation (so-called “dry” merging), but so far observational evidence is inconclusive and conflicting (e.g., van Dokkum 2005; Bell et al. 2006; Faber et al. 2005).

Understanding galaxy formation boils down to two questions: When did galaxies form their stars? And, when did they assemble into their present-day configurations? In these proceedings, I discuss constraints on star formation in massive galaxies at $z \sim 1.5-3$. In particular I focus on observations of the IR activity in high redshift galaxies using *Spitzer* 24 μm observations, and what this means for the galaxies’ assembly and evolution.

Star Formation in High- z Massive Galaxies

Deep *Spitzer* surveys at 24 μm show that IR luminous galaxies evolved very rapidly (Papovich et al. 2004), dominating the SFR density at $z \sim 1$ (e.g., Le Floc’h et al. 2005). Several studies of the 24 μm emission of higher redshift galaxies ($1.5 < z < 3$) show very high detection rates ($\gtrsim 50\%$; Daddi et al. 2005; Papovich et al. 2006; Reddy et al.




Figure 1. Specific SFR as a function of stellar mass for high redshift galaxies (from Papovich et al. 2006). The left panel shows galaxies at $1.5 < z < 3.0$. Red circles correspond to DRGs; gray triangles denote SFR upper limits. Blue squares show galaxies from the HDF-N. Stars show X-ray sources. The right panel shows galaxies from COMBO-17 at $0.65 < z < 0.75$. Dots denote SFR upper limits.

2006; Webb et al. 2006), suggesting that the majority of high-redshift galaxies emit in the thermal IR — *they are either actively forming stars, supermassive blackholes, or both at this epoch.*

For example, figure 1 shows the specific SFRs ($\Psi/\mathcal{M}$) derived from the masses and SFRs for the galaxies in the GOODS fields at $1.5 < z < 3.0$ (from Papovich et al. 2006), where the SFRs are derived from the summed UV and IR emission based on the *Spitzer* 24 μm data. The figure also shows the specific SFRs for $0.65 < z < 0.75$ galaxies from COMBO-17 (Wolf et al. 2003), derived also using *Spitzer* 24 μm data. Interestingly there is strong evolution in the specific SFRs, especially for massive galaxies. Galaxies with masses $\geq 10^{11} \mathcal{M}_{\odot}$ at $1.5 \leq z \leq 3$ have high specific SFRs, $\Psi/\mathcal{M} \sim 0.2\text{--}10 \text{ Gyr}^{-1}$, excluding X-ray sources. In contrast, at $z \lesssim 0.75$ galaxies with $\mathcal{M} \geq 10^{11} \mathcal{M}_{\odot}$ have much lower specific SFRs, $\Psi/\mathcal{M} \sim 0.1\text{--}1 \text{ Gyr}^{-1}$.

Papovich et al. (2006) defined the integrated specific SFR as the ratio of the sum of all galaxy SFRs, Ψ , to the sum of their stellar masses, $\mathcal{M}$. Figure 2 shows the integrated specific SFRs for galaxies selected from GOODS at $z \sim 1.5\text{--}3.0$ and COMBO-17 at $z \sim 0.4$ and 0.7 , all with $\mathcal{M} \geq 10^{11} \mathcal{M}_{\odot}$. The error box indicates the affect of assumptions in the SFRs and AGN activity (see further discussion below, and in Papovich et al. 2006). The integrated specific SFR in galaxies with $\mathcal{M} > 10^{11} \mathcal{M}_{\odot}$ declines by more than an order of magnitude from $z \sim 1.5\text{--}3$ to $z \lesssim 0.7$. The curves in figure 2 show the specific SFR integrated over all galaxies, not just the most massive; this is the ratio of the cosmic SFR density to its integral, $\dot{\rho}_* / \int \dot{\rho}_* dt$. Although there is a decrease in the global specific SFR with decreasing redshift, the evolution in the integrated specific SFR in massive galaxies is accelerated. The implication is that at $z \gtrsim 1.5$, massive galaxies are rapidly forming their stars, whereas by $z \lesssim 1.5$ the specific SFRs of massive galaxies drops rapidly, and lower-mass galaxies dominate the cosmic SFR density.

If AGN contribute to the observed 24 μm emission in galaxies at $z \sim 1.5\text{--}3$, then they can affect the inferred IR luminosities. For example, using an IR template for Mrk 231 instead of a star-forming galaxy with $L_{\text{IR}} \gtrsim 10^{13} L_{\odot}$ would reduce the IR luminosity for $z \sim 1.5\text{--}3$ galaxies by a factor of $\sim 2\text{--}5$. Many ($\sim 15\%$) of the massive galaxies at $z \sim 1.5\text{--}3$ are detected in the deep X-ray data (see Papovich et al. 2006), and these objects tend




Figure 2. The integrated specific SFR as a function of redshift (from Papovich et al. 2006). The integrated specific SFR is the ratio of the sum of galaxy SFRs to the sum of galaxy stellar masses. The curves show the expected evolution from the global SFR density. Data points show results for galaxies with $\geq 10^{11} M_\odot$. The inset error bar shows an estimate on the systematics.

to have high inferred specific SFRs (see figure 1). This obviously has an effect on the evolution of the integrated specific SFRs: the error box on the data point at $1.5 < z < 3$ in figure 2 shows how the result changes if the SFR for galaxies with putative AGN (based on X-ray detections, or rest-frame near-IR colors, see Stern et al. 2005; Alonso-Herrero et al. 2006) is set to zero. Interestingly, the high AGN occurrence in massive galaxies suggests that at $z \sim 1.5-3$ these objects are forming simultaneously their stars and supermassive black holes. This may provide the impetus for the present-day black-hole-bulge-mass relation and/or provide the feedback necessary to squelch star-formation in such galaxies, moving them onto the red sequence.

Confrontation with Models

Recent hierarchical galaxy-formation models predict a “downsizing” trend in the star formation rates of massive galaxies. De Lucia et al. (2006) show that within the semi-analytical galaxy-formation prescription coupled with feedback from AGN (Croton et al. 2006) that galaxies in the most massive dark matter haloes formed their stars at the earliest epochs. These models seem broadly consistent with observations. The “downsizing” jargon used by astronomers merely signifies that star-formation is accelerated in the most massive overdensities in current Λ CDM galaxy-formation models.

While encouraging, the details of star-formation in massive galaxies are not yet fully consistent with the *Spitzer* 24 μ m observations. Assuming the galaxies at $1.5 < z < 3$ with observed masses $> 10^{11} M_\odot$ evolve to present-day galaxies with masses of *at least* this much, then the De Lucia et al. model predicts they should have observed specific SFRs $\sim 0.3 \text{ Gyr}^{-1}$ on average. The observations instead suggest a specific SFR value closer to 1 Gyr^{-1} (see figure 1). There may be a discrepancy at the factor ~ 3 level, and this is likely discrepancy is likely greater because the observed galaxies presumably will continue to increase their stellar mass to $z \sim 0$. Admittedly the uncertainties of the IR-inferred SFRs are at the factor ~ 3 level, and to provide constraints on models of the star formation rates of massive galaxies will require lowering these observational uncertainties.

A more serious challenge to hierarchical models is the existence of a substantial population of apparently passive galaxies on the red sequence by $z \sim 1$. The stellar mass density of galaxies on the red sequence is near its present-day value by $z \sim 0.7$ (e.g., Brown et al. 2006). Furthermore, Cimatti et al. (2006; see also Cimatti et al., *these proceedings*) find that by using empirical color–evolution models the number of the most luminous galaxies ($>4 L^*$) on the red sequence is nearly unchanged to $z \sim 1$. This is difficult for hierarchical models that predict the most massive present-day galaxies assembled into their current configurations at the lowest redshifts (e.g., Neistien et al. 2006; De Lucia et al. 2006). For example, both the Neistien et al. and De Lucia et al. show that the main progenitor (having $\gtrsim 50\%$ of the final mass) of massive galaxies with present-day $M > 10^{12} M_\odot$ formed at $z < 1$. This is difficult to reconcile even with frequent “dry merging” on the red sequence (see also Faber et al. 2005).

We are witnessing a growing understanding (and even possible convergence) between observational and theoretical constraints on galaxy evolution, even though significant hurdles remain. It will be exciting to see the summary of our knowledge in this field at the next IAU General Assembly.

Acknowledgements

I wish to thank the IAU symposium organizers for the invitation to present this material, and for planning a successful meeting in such a historic locale. I am grateful for my colleagues on the MIPS GTO and GOODS teams for their continued collaboration. Support for this work was provided by NASA through the Spitzer Space Telescope Fellowship Program, through a contract issued by JPL/Caltech under a contract with NASA.

References

- Alonso-Herrero, A., et al. 2006, *Astrophys. J.*, 640, 167
- Bell, E. F., Wolf, C., Meisenheimer, K., et al. 2004, *Astrophys. J.*, 608, 752
- Bell, E. F., et al. 2006, *Astrophys. J.*, 640, 241
- Blanton, M. R., Hogg, D. W., Bahcall, N. A., et al. 2003, *Astrophys. J.*, 594, 186
- Chary, R. R., & Elbaz, D. 2001, *Astrophys. J.*, 556, 562
- Cimatti, A., Daddi, E., & Renzini, A. 2006, aap, 453, 29
- Croton, D., Springel, V., White, S. D. M., et al. 2006, *Mon. Not. Royal. Astro. Soc.*, 365, 11
- Daddi, E. et al. 2005b, *Astrophys. J. Lett.*, 631, L13
- Davé, R. et al. 2005, preprint (astro-ph/0510625)
- Dickinson, M., Papovich, C., Ferguson, H. C., & Budavári, T. 2003, *Astrophys. J.*, 587, 25
- Faber, S., et al. (2006), preprint (astro-ph/050644)
- Hopkins, P., et al. 2006, *Astrophys. J. Sug.*, 163, 1
- Juneau, S., Glazebrook, K., Crampton, D., et al. 2005, *Astrophys. J. Lett.*, 619, L135
- Kauffmann, G., & Charlot, S. 1998, *Mon. Not. Royal. Astro. Soc.*, 294, 705
- Le Floc’h, E., Papovich, C., Dole, H., et al. 2005, *Astrophys. J.*, 632, 169
- McCarthy, P. 2004, *Ann. Rev. Astron. & Astrophys.*, 42, 477
- Papovich, C., Dole, H., Egami E., et al. 2004, *Astrophys. J. Sug.*, 154, 70
- Papovich, C., Moustakas, L. A., Dickinson, M., et al. 2006, *Astrophys. J.*, 640, 92
- Reddy, N. et al. 2006, *Astrophys. J.*, 644, 792
- Renzini, A. 2006, *Ann. Rev. Astron. & Astrophys.*, 44, 141
- Rudnick, G., et al. 2006, *Astrophys. J.*, 650, 624
- Stern, D., Eisenhardt, P., Gorjian, V., et al. 2005, *Astrophys. J.*, 631, 163
- Webb, T. M. A. et al. 2006, *Astrophys. J.*, 616, L17
- Willmer, C. N. A., et al. (2006), *Astrophys. J.*, 647, 853
- Wolf, C. et al. 2003, *Astron. & Astrophys.*, 401, 73
- van Dokkum, P. 2005, *Astronom. J.*, 130, 264